

Two remarks on wetting and emulsions

P. G. de Gennes
Collège de France, 11 place M. Berthelot,
75231 Paris Cedex, France

February 1, 2008

Abstract

This paper is extracted from an opening address given at the workshop "Wetting: from microscopic origins to industrial applications" (Giens, May 6-12, 2000). It discusses two special points a) the nature of line energies for a contact line b) the aging of emulsions.

1 Introduction

These notes are dedicated to the memory of Karol Mysels. I met him only rather late (in the 1980's). But I learned a large number of important things from him. I also treasure a certain group of his original slides on soap films, which he gave to me, and which I have used in a number (~ 200) of talks in high schools. Karol was both a gifted experimentalist and a deep thinker. It was one of my great prides to have Karol and Estella as participants for the Stockholm ceremonies, in which I was involved in 1991.


The following notes are a (clumsy) attempt to contribute to some things in which Karol was deeply interested: the basics of wetting, and the stability of foams and emulsions. The notes are rough, and would have been much improved if he had still been with us... We shall not forget him.

2 Line energies

If we pinch a violin string (fig. 1), we obtain a static triangular form. This is associated with a standard line tension $\mathfrak{S}$. For small displacements u with Fourier transforms u_q the energy has the standard form:


$$E = \sum_q \frac{1}{2} \mathfrak{S} q^2 |u_q|^2 \quad (1)$$

On the other hand, if we pinch a contact line (for instance between a fluid, a solid, and air) we find a very different form (fig. 2). This originates from a different structure of the energy:



$$E = \sum_q \frac{1}{2} \gamma f(\theta_e) |q| |u_q|^2 \quad (2)$$

where θ_e is the equilibrium contact angle, γ is the surface tension, and $f(\theta_e)$ is a dimensionless function, discussed in refs [1] and [2]. The $|q|$ dependence is very singular: it comes from the integration of q^2 contributions over the width of the perturbed fluid region (q^{-1}).



Eq. 2 corresponds to what I call "fringe elasticity". It holds for scales q^{-1} which are smaller than the Laplace length:

$$\varkappa^{-1} = \left(\frac{\gamma}{\rho g} \right)^{1/2} \quad (3)$$

ρ : density, g : gravitational acceleration.

It is well known that we should add in equation (2) an intrinsic line tension $\mathfrak{S}_0$, transforming eq. (2) into:

$$E = \sum_q \left\{ \frac{1}{2} \gamma f(\theta_e) |q| + \mathfrak{S}_0 q^2 \right\} \quad (4)$$

Molecular models give values of $\mathfrak{S}_0$ (positive or negative) which are of order:

$$\mathfrak{S}_0 = \gamma a \quad (5)$$

where a is a molecular size (a few angstroms).


At submillimeter scales ($a < q^{-1} < \kappa^{-1}$) this correction should be unobservable. However, a number of groups [3] [4] [5] [6] [7] have tried to measure $\mathfrak{S}_0$ by looking at the contact angle in small droplets. The result can be stated in the form:

$$\mathfrak{S}_0 = \gamma \ell \quad (6)$$

where $|\ell|$ is often anomalously large (of order 1 micron).

My own view is that this is an artefact of optical methods. One example, based on ray optics, is illustrated on fig. 3. In principle, we determine θ through the maximum deflection angle of a reflected ray $\varphi = 2\theta_e$. However, because of diffraction effects, the actual last measurement point is at a distance $\sim \lambda$ from the contact line, and has a weaker deflection $\varphi - \epsilon$, where $\epsilon \sim \lambda/R$ (R : radius of curvature; λ : optical wavelength). This correction has exactly the same structure as the inclusion of $\mathfrak{S}_0$, and gives an apparent $\mathfrak{S}_0$:

$$\mathfrak{S}_0/app \sim \gamma \lambda \quad (7)$$



This is the artefact. I believe that all optical methods have the same defect.

3 Aging of emulsions

Certain O/W emulsions, which are monodisperse, can coalesce slowly, with droplets growing in size, but remaining monodisperse [8]. This occurs in the absence of contaminants. In practice, many practical emulsions do contain some small particles, with a wettability such that the particle gets hooked at the O/W interface, although it is rather hydrophobic. These grains induce the rupture of oil films by the so called Garrett process [9]. If the grains are very dilute (so that each grain works individually), this can lead to a rough, polydisperse emulsion. Our aim here, is to discuss this, at the level of scaling laws, following the lines of ref .[10].

Start with a single grain: when trapped on a film, it induces rupture within a certain time τ . When the film is locally destroyed, we postulate that the particle binds to a neighboring interface -the size of the jump being comparable to the diameter of the initial droplets.

We are thus led to think first of a random walk with a diffusion coefficient:

$$D = d^2/\tau \quad (8)$$

This walk destroys some droplets: the overall amount of interface present decreases, and this could alter the control parameters: we want to avoid this to reach a simple discussion:

a) We assume that there is a plentiful reservoir of surfactant (concentration above *cmc*).

b) The thickness of the water films could increase, and this would reduce the coalescence rates. If we estimate the diffusion coefficient D_W of *water* in the structure, we find that it is related to Poiseuille films in the Plateau borders (of diameter h) induced by gradients of the pressure:

$$p \cong p_0 - \gamma/h \quad (9)$$

The (qualitative) result is:


$$D_W \cong \frac{\gamma}{\eta_W} h \quad (10)$$

where η is the water viscosity. In all what follows, we assume $D_W \gg D$. Then water can diffuse fast from the "wounded" region to the bulk of the monodisperse emulsion. It is then plausible to assume that the water films remain at constant thickness.

A major point is the following: the grain does *not* follow an arbitrary random walk in the emulsion. If it "digs a tunnel" in the network of droplets, the Laplace pressure at the bottom of the tunnel forces it to retract very soon.

This retraction would follow the Washburn law for fluid motion in a capillary, with a certain diffusion coefficient D_r (r stands for retraction). We assume (as in agreement with usual conditions) that $D_r > D_W > D$.

In this situation, the grain can move only at the interface between a coarse oil drop and the unperturbed fine emulsion, as shown on fig. 4.



It builds up a coarse drop of radius $R(t)$ increasing with time t . The number of ruptured droplets is $t/\tau (>> 1)$, and the oil volume of the coarse drop is (omitting coefficients):

$$R^3 = \frac{t}{\tau} d^3 = dDt \quad (11)$$

During time t , the grain has moved at random on the surface of the coarse drop, spanning a length s such that $s^2 = Dt$. We see from this that:

$$\frac{\ell^2}{R^2} = \frac{R}{d} >> 1 \quad (12)$$

This means that the particle explores fast the surface, and ensures that the coarse droplet is essentially spherical. (For more details on the roughness of the interface, see ref.[10].

Our conclusions are the following:

If we start, at $t = 0$, with a monodisperse emulsion, containing a very dilute suspension of active grains, we expect a first stage, where each grain generates one coarse droplet of size $R(t)$, growing like $t^{1/3}$.

At a certain moment ($t = t^*$), these coarse droplets enter into contact. At $t > t^*$, the original fine emulsion is not the main component, and the growth laws must become more complex. Even if D_W is fast, the water films will thicken, and coalescences should slow down, or even stop, as often observed. On the other hand, if τ has become long, one could possibly arrive at a fusion of the coarse droplets, leading to an inverted (W/O) emulsion.

On the whole, we are still very far from understanding the aging of emulsions. But we clearly have an intrinsic blow up of the walls (which here are water films) plus an extrinsic process due to grains (or other external objects). Monodisperse emulsions may allow us to separate neatly the two processes.

References

- [1] J. F. Joanny, P. G. de Gennes, J. Chem. Phys. **81**, 552-562 (1984).
- [2] F. Brochard-Wyart, D. Quéré, P. G. de Gennes "Gouttes, bulles et perles", Eyrolles ed. (to be published).
- [3] D. Li, A. Neuman, Colloids Surf., **43**, 195 (1990).
- [4] J. Drelich, J. Wilbur, J. D. Miller, G. Whitesides, Langmuir, **12**, 1913 (1996).
- [5] J. Drelich, Polymer J. (Chem), **71**, 525 (1997).
- [6] A. Amirfazli, D. Kwok, J. Gaydos, A. Neuman, J. Colloid Interface Sci., **205**, 1 (1998).
- [7] J. Y. Wang, S. Betelu, B. M. Law, Phys. Rev. Lett., **83**, 3677 (1999).
- [8] B. Deminière, A. Colin, F. Leal Calderon, J. Bibette, Phys. Rev. Lett., **82**, n° 1, 229-232 (1999).
- [9] P. R. Garrett "Defoaming", Surfactant Science Series, **45** (M. Dekker, NY, 1997).
- [10] P. G. de Gennes, C. R. Acad. Sci. (Paris), **328 II**, 331-335 (1998).